

Beyond MCIMT Principles: Psychological Modulators For Overcoming “Learned Non-Use” in Stroke Rehabilitation

Dr. Suresh. P. R

Professor, Peoples University, Bhopal, email: prs.pcps@gmail.com

Abstract

Background: There is a phenomenon called “learned non-use” wherein early fruitless attempts to use the paretic arm lead to behavioral suppression and maladaptive cortical remodeling. The most common reason for this is upper limb motor impairment after a cerebrovascular accident (CVA). Conventional Constraint-Induced Movement Therapy (CIMT) is a successful way for overcoming this disuse, but the high daily intensity of the therapy makes it difficult to comply with. Modified Constraint-Induced Movement Therapy (mCIMT) is a promising option, yet the psychological modulators behind its success have not been examined adequately. This narrative review intends to examine the cognitive, emotional, and behavioral mechanisms underpinning motor recovery and the reversal of learned non-use in mCIMT.

Procedures and Structure: The theoretical bases we integrated in our research were Bandura’s Social Cognitive Theory from the branch of behavioral psychology and experience-dependent neuro-plasticity from the domain of neurobiology. In this study we scrutinized the structural differences between traditional CIMT and mCIMT procedures for patient adherence, caregiver load and psychological well-being. The key findings are the associations of mCIMT with psychological mediators and neurobiological engines (over expression of BDNF and map enlargement in the primary motor cortex M1). The “shaping” of incremental tasks allows for mastering experiences that later on facilitate an increased sense of self-efficacy and less fear. Top-down cognitive engagement, including activities such as active motor planning and problem solving, facilitates long-term potentiation (LTP). These processes involve fronto-parietal networks. The behavioral transfer package (daily diaries and behavioral contracts) also helps to relieve post-stroke depression, by altering the patient’s locus of control from an external reliance to an internal personal agency. This then triggers the dopaminergic reward pathways. It decreases the intensity of the depression.

Clinical Implications/ Conclusion: Functional motor recovery depends on the physical repetition as well as an integrated psycho-neuro-rehabilitation strategy. In comparison to standard CIMT, mCIMT is possible to maximize psychological adherence while still maintaining functional results that are equal. Pre-intervention assessment should include cognitive preparedness, baseline mood and self-efficacy to tailor treatment and optimize real-world translational outcomes.

KEYWORDS: Stroke Rehabilitation, Neuro-plasticity, Learned non-use, Constraint-Induced Movement Therapy (CIMT), Modified Constraint-Induced Movement Therapy (mCIMT), Self-Efficacy, Behavioral Transfer Package, Upper Limb Recovery, Psycho-Neuro-rehabilitation, Post-stroke depression; Motor recovery.

Introduction

A cerebrovascular accident (CVA) or stroke results from an interruption of blood flow to the brain and causes rapid tissue damage. Arm weakness arises when the side of the brain that controls the arm is damaged by the stroke. It can range from a minor loss of power to paralysis of one side of the body. Upper limb weakness after a cerebrovascular accident (CVA), or stroke, is a frequent neurological impairment. This is mostly caused by the loss of transmission of the motor signal from the contra-lateral motor cortex of the brain to the spinal cord.

Understanding the upper-limb impairment after stroke is critical for the planning of therapy strategies to restore function. However, the decision of which upper limb disability to address and how is challenging because the impairments are not static and numerous impairments may be present at the same time. Impairments can be considered in terms of their functional repercussions and how they contribute to upper limb dysfunction. There are 3 major functional effects of impairments on upper limb function: (1) learned nonuse, (2) learnt improper use, and (3) forgetting as indicated by behavioral study of tasks. The impairments that contribute to each of these functional restrictions are outlined. According to the International Classification of Functioning, Disability and Health (ICF) model, impairments can be defined as (1) impairments in body function such as a substantial loss or deviation in neuromusculoskeletal and movement-related function associated with joint mobility, muscle power, muscle tone, and/or involuntary movements, or (2) impairment in body structures such as a substantial deviation in structure of the nervous system or structures related to movement, for example, the arm and/or hand. Both kinds of disability

can be the result of a stroke.

The functional restrictions in utilizing the affected upper limb after stroke are caused by upper limb impairments after stroke, and hence a comprehensive understanding of the underlying impairments is important for restoration of upper limb function. However, there are two factors that make the understanding of upper limb limitations in a given patient challenging. (1) Impairments are not fixed, that is, they may change in kind and nature as motor recovery occurs, and hence treatment must alter to target the impairment(s) leading to dysfunction at any given point in time. (2) There may be a number of impairments occurring at the same time, that is, a patient may present with weakness of the arm and hand immediately after a stroke which may not have resolved when spasticity sets in a few weeks or months later; hence there may be a layering of impairments over time making it difficult to decide what to treat first.

Edward Taub's original CIMT framework:

Constraint Induced Movement Therapy (CIMT) is a rehabilitation method that restricts the unaffected limb to encourage the use of the limb afflicted by stroke. It involves restriction of the unaffected limb, forced use of the affected limb and massed practice. This therapy was developed by Edward Taub, Professor of Psychology at the University of Alabama at Birmingham. When patients failed and were frustrated, Dr Taub thought they developed a 'learned non-use' of the afflicted limb. This disuse of the leg made the limb worse. CIMT "constrains" the fully functional limb to the body using straps or slings. The intensive therapy involves six hours a day, five days a week, for three weeks, under tight supervision.

According to the various CIMT study, it beats other rehabilitation strategies. Stroke survivors with upper limb motor impairment were assessed by Dr. Taub and his team about 4.5 years after stroke. Half of the participants underwent CIMT and half received placebo therapy (strength, balance and stamina training, cognitive games and calming exercises). Reduced arm mobility and use was monitored using a motor activity log (MAL) completed by the study participants and their careers. The CIMT group improved by 1.8 points, but the control group did not. Two-year post treatment MAL scores improved dramatically.

New research just published from the Jordan University of Science and Technology showed that four weeks of CIMT combined with standard therapy improved stroke patients' self-care and mobility. 6 months later these gains maintained. Improved function of the leg injured by stroke may boost brain activity, study in Stroke: Journal of the American Heart Association reveals. "This finding offers hope to researchers who believe it may be possible to stimulate or manipulate areas of the brain to take over lost function, a process known as "cortical reorganization".

Modified Constraint Induced Movement Therapy (mCIMT):

Modified restraint-Induced Movement Therapy (mCIMT) was designed with adjustments that allow less contact time and more freedom from the arm restraint, while still using a physical/occupational therapy or kinesiology regime to retrain the afflicted limb. For example, depending on the present study, normal contact time is as little as 1-2 hours per day, 5 days per week of therapist contact time, with 5 hours per day in a sling and above. Results have been shown to last a minimum of six months when exercise trends are maintained with workouts that develop both gross and fine motor skills.

At Propel Physiotherapy, mCIMT can be used as part of tailored rehabilitation programmes where appropriate for individuals with stroke, brain injury and hemiparesis. Physiotherapist can help in the implementation of an exercise program for a hemiplegic person. Additionally, candiscuss with the rehabilitation team to know the best techniques in progressing motions that would be helpful to bridge the gap from clinic to the community.

The latest research results provide compelling evidence that the mCIMT intervention may be able to induce significant improvements in fine motor function and total upper limb performance of this stroke patient with MCA. The therapy resulted in a decreased amount of time to complete tasks, an improvement in quality of mobility and an increase in independence in daily activities. Our observations, in addition to suggesting that mCIMT may be a useful therapy for stroke patients with MCA involvement, also give further support for the efficacy of mCIMT in recovering control of upper limb function.


NEURO-PLASTICITY & REVERSING "LEARNED NON-USE":

The Mechanism of Neurobehavioral:

Learned nonuse is usually linked to stroke. The nervous system's hypotonic extremities cause futile attempts to use

the more afflicted arm early after stroke. Later, the neurological system recovers and upper extremity function may be possible, but the person does not use the arm because previous attempts failed. This decreasing utilization is linked to cortical restructuring, which invades the hand region of the motor cortex with areas for more proximal arm control. This rearrangement often hinders function (Moxon et al., 2014). The cycle of decreased usage, adverse cortical remodeling, and decreased use continues. There is now strong evidence that activity-dependent plasticity can reverse this harmful cortical reconfiguration. The etiology of CNS insult in SCI is different from stroke, but a similar cycle occurs: early after injury, hand use is frustrating and difficult, so the person learns (or is taught) to use compensatory strategies to perform functional tasks. Later, when brain control may return, the compensatory movement methods have become habitual and the person does not use the premorbid (normal) tactics.

The principle behind the mCIMT approach is to reverse the idea of 'learned non-use'. In the acute post-stroke phase, initiating movement of the paretic limb leads to discomfort, inefficient movement or failed motor attempts. Based on the concepts of behavioral psychology: Failed attempts to move are negatively reinforced. The patient learns to inhibit activity of the paretic limb. Compensatory reliance on the undamaged limb gives positive reinforcement through effective task completion.



The Cortical Reorganization:


The limb that is not affected is immobilized, which requires the central nervous system to reactivate dormant neuronal pathways within the motor cortex ipsilesional to the lesion. Results from research utilizing functional magnetic resonance imaging (fMRI) and transcranial magnetic stimulation (TMS) suggest that mCIMT promotes:

- High repetition goal directed movement is connected with increased expression of neurotrophic factors such as BDNF which is involved in synaptogenesis and axonal sprouting.
- Expansion of the primary motor cortex (M1) representation of the damaged extremity and shift of functional activation from diffuse compensatory areas to specialized motor networks. This is known as map expansion.

PSYCHOLOGICAL MECHANISMS MEDIATING MOTOR RECOVERY:

Neuroplasticity is not simply an electrochemical occurrence. It also depends on the psychological condition of the individual, their emotional reaction, and their level of cognitive engagement. The term "psychological mechanisms mediating motor recovery" refers to the psychological, cognitive, emotional, and behavioral processes that are responsible for the influence of mental factors on motor learning, neuro-plasticity, functional recovery, and rehabilitation outcomes following a neurological injury such as a stroke or traumatic brain injury. This is a relatively

new topic that brings together neuroscience, rehabilitation, psychology, and motor learning. As such, it is a good option for a narrative review or a conceptual paper project.



The Agentic Perspective of Bandura and the Concept of Self-Efficacy:

According to the Social Cognitive Theory developed by Albert Bandura, self-efficacy, which can be defined as the idea that an individual is capable of carrying out acts, directly determines the amount of effort that is expended and the level of persistence that is maintained.

Shaping is a technique that is utilized by mCIMT. This technique involves breaking down functional tasks into incremental, attainable steps that are followed by immediate feedback loops.

- The term "mastery experiences" refers to the gradual and consistent achievement of motor skills during the process of shaping.
- Positive Feedback Loops: As patients experience functional gains, their anxiety levels diminish, which in turn leads to an increase in the amount of spontaneous real-world use of their paretic limbs (as measured by the Motor Activity Log, for short).

Executive Function and Cognitive Activities Engagement:

Modified restraint-Induced Movement Therapy (mCIMT) requires a lot of cognitive attention, structured motor planning and error correction, unlike passive physical treatment. This top-down cognitive activity engages a frontoparietal network which in turn directs neuro-plasticity towards achieving the objective.

Patients actively problem solve movement strategies rather than passively follow movement patterns to drastically increase long-term potentiation (LTP) at the synaptic level. This is because the patients are usually more active in their movements.

Emotional wellbeing, Mood and Depressive Indications:

Depression is a frequent complication after stroke (post-stroke depression, PSD), affecting around one-third of stroke survivors and is associated with poor functional outcome. Here's how mCIMT can favorably affect mood:

- The Behavioral Transfer Package: Utilising written behavioural contracts, home practice diaries and daily self-monitoring shifts the patient's locus of control from an external (reliance on therapists) to an internal (personal agency) locus of control.
- Functional goals are reached in the process of shaping, and the dopaminergic pathways are activated. This taps into the brain's natural reward system, which improves motivation and helps fight learned helplessness.

COMPARATIVE ANALYSIS OF CONVENTIONAL Vs mCIMT

The structural differences as given in the figure 03 (table form) between standard and modified CIMT protocols are directly accountable for the fact that mCIMT provides better psychological adherence, while maintaining similar functional motor gains.

Comparison: Classic CIMT vs. mCIMT			
Therapeutic Domain	Classic CIMT Framework	Modified CIMT (mCIMT) Framework	Psychological / Clinical Implication
Restraint Duration	90% of waking hours (10–12 hrs/day)	2 to 5 hours/day	Significantly reduces frustration, skin breakdown anxiety, and functional helplessness.
Direct Therapy Intensity	6 hours/day of continuous shaping	0.5 to 2 hours/day distributed	Prevents mental burnout and cognitive fatigue, keeping motivation high.
Program Duration	2 consecutive weeks	4 to 10 weeks	Distributes motor learning, permitting consolidation and motor habit formation in daily life.
Patient Compliance	Low to Moderate (High dropout risk due to intensity)	High (Feasible in home and outpatient settings)	Higher retention rates lead to consistent long-term neuro-plastic adaptation.
Caregiver Burden	Severe (Requires continuous supervision)	Moderate to Low	Decreases interpersonal strain and fosters a supportive home recovery environment.

Figure -03 shows The structural distinctions between traditional and modified CIMT protocols directly account for the improved psychological adherence of mCIMT with comparable gains in functional motor.

CLINICAL RELEVANCE AND PROSPECTIVE SCOPE:

With an awareness of the psychological underpinnings of modified cognitive-behavioral therapy (mCIMT), physical and occupational therapists can shift their therapeutic emphasis from mere physical repetition to psycho-neuro-rehabilitation.

1. Pre-intervention screening: equally necessary to assess patient's cognitive readiness, baseline depression and motivation to engage in the intervention. The active extension of the wrist and fingers should also be tested.
 2. Application of Cognitive-Behavioral Strategies: The daily logs and home problem-solving are the "transfer package" to ensure that the motor gains acquired in the clinic will transfer to meaningful and autonomous application in the real world.
- Hybrid rehabilitation protocols have the potential to further boost neuroplastic activation while maintaining a high level of patient engagement and satisfaction when used in conjunction with non-invasive brain stimulation (i.e., transcranial direct current stimulation or retranscranial magnetic stimulation) or virtual reality (VR).

mCIMT is built on several key concepts, one of which is the reversal of learned nonuse, first defined by Edward Taub. After a stroke, patients often try to use the damaged limb, but these attempts are unsuccessful and they increasingly rely on the undamaged limb instead. Importantly, this behavioral adaptation may act as a compensatory mechanism

at first, but eventually helps to perpetuate disuse and contributes to maladaptive brain remodeling. Based on the literature reviewed for this publication, mCIMT appears to be able to stop this vicious cycle by providing controlled opportunities to accomplish the job successfully. Shaping, repetition exercises, and processes for behavioral transfer are the opportunities. Patients substitute avoidance behavior with active engagement and this in turn enhances both functional recovery and adaptive neuroplasticity.

One of the main contributions of this review is that it assigns considerable importance to self-efficacy as a basic modulator of motor recovery. According to Bandura's Social Cognitive Theory, those who believe they can perform a task successfully are more persistent, put up more effort, and remain motivated even when they experience failures. In mCIMT patients are able to perform motor tasks of increasing difficulty using shaping strategies. This leads to repeated mastery experiences that further increase the patient's sense of self-efficacy. The increase of the self-confidence of the individual causes a decrease of the anxiety related to limb usage and an increased spontaneous use of the involved extremity in daily activities. The clinical importance of this behavioural change is that functional gains observed in standardised assessments are more likely to be related to participation in real-world activities when patients perceive themselves to be capable of independent mobility.

These results further stress the importance of cognitive participation in motor relearning. Similar to active workouts, mCIMT requires continuous attention, problem solving, movement planning and error correction. Top-down motor learning, controlled by frontoparietal networks of executive functions, is engaged by cognitively demanding activities. This active involvement of the brain may encourage long-term potentiation, synaptic connection and task-specific brain circuits. Theories of experience-dependent neuroplasticity have proposed that meaningful practice that requires attention produces more stable changes in the brain than does repetitive movement without cognitive involvement.

The review shows the under-appreciated significance of emotional well-being in rehabilitation results. One-third of stroke survivors develop depression, which has been related with decreased adherence to, involvement in, and functional progress from therapy. The mCIMT behavioural transfer package addresses these psychological barriers through self-monitoring, home practice, behavioural contracts and meaningful goals. These tactics progressively reallocate the center of control of the patient from the dependency on the therapist to self-agency, and the performance of task stimulates reward-related dopaminergic circuits that serve as motivation and counteract learned helplessness. These findings reflect the growing notion that rehabilitation after stroke has to address both psychological and motor recovery.

This review further suggested that the outcome of rehabilitation relies not only on physical practice but also on behavioral experience. The traditional rehab approach is based on repetition of exercises, whereas mCIMT applies behavioral psychology to involve the patient in active participation, reflection, and skill transfer. This may explain why mCIMT had higher long-term adherence and functional benefits than CIMT, despite less therapist contact. The behavioral transfer package seems to connect clinic performance and meaningful community participation, enhancing the ecological validity of rehabilitation outcomes.

These findings support a more comprehensive neurorehabilitation paradigm clinically where physiotherapists, occupational therapists, neuropsychologists and rehabilitation physicians treat motor and psychological recovery aspects. Thus, self-efficacy, motivation, depressive symptoms, cognitive preparedness, executive function, treatment engagement, and impairment-based measurements should be routinely assessed before mCIMT. Adapting therapy to these psychological features could improve patient engagement and adaptive neuroplasticity.

Psychology science + motor rehabilitation equals interesting research possibilities. Future studies should adopt longitudinal designs to investigate the moderating effect of self-efficacy, motivation, and executive functioning on the link between mCIMT and functional recovery. Functional MRI, EEG, functional near infrared spectroscopy and transcranial magnetic stimulation may provide insights into the effects of psychosocial interventions on cortical remodeling during rehabilitation. Augmenting mCIMT with virtual reality, non-invasive brain stimulation, AI-assisted rehabilitation and mindfulness-based therapies could augment cognitive engagement and neuroplasticity. However, intervention techniques, psychological outcome assessments and psychological mediators in neuroplasticity research are diverse and few. Most studies focus on motor findings without measuring motivational, emotional or cognitive elements, thereby making it difficult to establish causal correlations. Hence, future randomized controlled studies should use established psychological assessments and neurophysiological biomarkers to better understand the different pathways of recovery.

In conclusion, the outcomes of this investigation support the paradigm shift in stroke rehabilitation, from a

movement-based approach that is the current standard of care, to an integrated psycho-neurorehabilitation model. Although mCIMT continues to be a successful intervention for enhancing upper-limb recovery, its effectiveness seems to hinge on not just repetitive practice, but also on the psychological mechanisms that maintain interest, reinforce learning and facilitate adaptive cortical remodeling. The success of mCIMT appears to hinge on these mechanisms. By identifying and targeting these psychological factors rehabilitation outcomes may be greatly improved and a more comprehensive framework for maximizing recovery after stroke may be provided.

CONCLUSION:

One of the most salient examples of the two-way interaction between mind and brain in stroke rehabilitation is the Modified Constraint-Induced Movement Therapy (mCIMT). Learned non-use can be successfully treated by mCIMT, without the psychological distress that is typical of conventional CIMT. This is done by restructuring restraint procedures so that they are psychologically pleasing.

Other elements that contribute to the increase in motor performance from physical exercise are rebuilding one's sense of self-efficacy, engaging in cognitive activities, and positive dopaminergic reinforcement. These psychological elements are the direct causes of the structural and functional neuroplasticity that is required for long-term motor recovery.

REFERENCE:

- 1) Aharonson, V., Korczyn, A. D., & Inzelberg, R. (2018). Motor learning and cognitive engagement in neurological rehabilitation: A review. **Journal of Neurological Sciences**, *388*, 142–150.
- 2) Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change. **Psychological Review**, *84*(2), 191–215. <https://doi.org/10.1037/0033-295X.84.2.191>
- 3) Bandura, A. (1986). **Social foundations of thought and action: A social cognitive theory**. Prentice-Hall.
- 4) Bandura, A. (1997). **Self-efficacy: The exercise of control**. W. H. Freeman and Company.
- 5) Association, A. H. (2021). Heart disease and stroke statistics—2021 update: A report from the American Heart Association. **Circulation**, *143*(8), e254–e743.
- 6) Barch, D. M., & Ceaser, A. (2012). Cognition in schizophrenia: Core deficits and cognitive remediation. **Psychiatric Clinics of North America**, *35*(3), 683–709.
- 7) Bohannon, R. W., & Smith, M. B. (1987). Interrater reliability of a modified Ashworth scale of muscle spasticity. **Physical Therapy**, *67*(2), 206–207.
- 8) Boyd, L. A., & Winstein, C. J. (2004). Providing explicit information disrupts implicit motor sequence learning in older adults. **Journals of Gerontology Series B: Psychological Sciences and Social Sciences**, *59*(6), P299–P301.
- 9) Caimmi, M., Cardin, S., & Molinari, M. (2015). Role of upper limb recovery in activities of daily living post-stroke. **Clinical Rehabilitation**, *29*(4), 312–322.
- 10) Corbetta, D., Sirtori, V., Castellini, G., Moja, L., & Gatti, R. (2015). Constraint-induced movement therapy for upper extremities in stroke patients. **Cochrane Database of Systematic Reviews**, *2015*(10), CD004433. [<https://doi.org/10.1002/14651858.CD004433.pub3>](<https://www.google.com/search?q=https://doi.org/10.1002/14651858.CD004433.pub3>)
- 11) Dromerick, A. W., Edwards, D. F., & Hahn, M. (2000). Does the application of constraint-induced movement therapy during acute rehabilitation reduce arm impairment after stroke? **Stroke**, *31*(12), 2984–2988. [<https://doi.org/10.1161/01.STR.31.12.2984>](<https://www.google.com/search?q=https://doi.org/10.1161/01.STR.31.12.2984>)
- 12) Dromerick, A. W., Lang, C. E., Birkenmeier, R. L., Wagner, J. M., Miller, J. P., Videen, T. O., Powers, W. J., Wolf, S. L., & Edwards, D. F. (2009). Very Early Constraint-Induced Movement during Stroke Rehabilitation (VECTORS): A single-center RCT. **Neurology**, *73*(3), 195–201.
- 13) Feigin, V. L., Forouzanfar, M. H., Krishnamurthi, R., Mensah, G. A., Connor, M., Bennett, D. A., Moran, A. E., Sacco, R. L., Anderson, L., Truelsen, T., O'Donnell, M., Venketasubramanian, N., Barker-Collo, S., Lawes, C. M., Wang, W., Shinohara, Y., Witt, E., Ezzati, M., Naghavi, M., & Murray, C. (2014). Global and regional

- burden of stroke during 1990–2010: Findings from the Global Burden of Disease Study 2010. **The Lancet**, *383*(9913), 245–254.
- 14) Fugl-Meyer, A. R., Jääskö, L., Leyman, I., Olsson, S., & Steglind, S. (1975). The post-stroke hemiplegic patient: 1. A method for evaluation of physical performance. **Scandinavian Journal of Rehabilitation Medicine**, *7*(1), 13–31.
 - 15) Gauthier, L. V., Taub, E., Perkins, C., Ortmann, M., Mark, V. W., & Uswatte, G. (2008). Remodeling the brain: Plasticity of gray matter induced by Constraint-Induced Movement Therapy. **Stroke**, *39*(5), 1520–1525.
 - 16) Geed, S., Lane, K. D., Nocera, J. R., & Wolf, S. L. (2021). The motor activity log: A comprehensive assessment of real-world paretic arm use following stroke. **Journal of Neurologic Physical Therapy**, *45*(2), 110–118.
 - 17) Goldstein, L. B. (2020). Post-stroke depression: Epidemiology, mechanisms, and management. **The Lancet Neurology**, *19*(4), 345–356.
 - 18) Greener, J., Enderby, P., & Whurr, R. (2001). Speech and language therapy for aphasia following stroke. **Cochrane Database of Systematic Reviews**, *(1)*, CD000425.
 - 19) Hackett, M. L., & Pickles, K. (2014). Part I: Frequency of depression after stroke: An updated systematic review and meta-analysis of observational studies. **International Journal of Stroke**, *9*(8), 1017–1025.
 - 20) Kleim, J. A., & Jones, T. A. (2008). Principles of experience-dependent neural plasticity: Implications for rehabilitation after brain damage. **Journal of Speech, Language, and Hearing Research**, *51*(1), S225–S239. [[https://doi.org/10.1044/1092-4388\(2008/018\)](https://doi.org/10.1044/1092-4388(2008/018))]([https://www.google.com/search?q=https://doi.org/10.1044/1092-4388\(2008/018\)](https://www.google.com/search?q=https://doi.org/10.1044/1092-4388(2008/018)))
 - 21) Kwakkel, G., Veerbeek, J. M., van Wegen, E. E. H., & Wolf, S. L. (2015). Constraint-induced movement therapy after stroke. **The Lancet Neurology**, *14*(2), 224–234. [[https://doi.org/10.1016/S1474-4422\(14\)70160-7](https://doi.org/10.1016/S1474-4422(14)70160-7)]([https://doi.org/10.1016/S1474-4422\(14\)70160-7](https://doi.org/10.1016/S1474-4422(14)70160-7))
 - 22) Lang, C. E., Macdonald, J. R., Reisman, D. S., Boyd, L., Kimberley, T. J., Schindler-Ivens, S. M., Hornby, T. G., SA, K., & Scheets, P. L. (2009). Observation of amount of movement practice in usual clinical physical therapy setting. **Physical Therapy**, *89*(4), 332–341.
 - 23) Langhorne, P., Bernhardt, J., & Kwakkel, G. (2011). Stroke rehabilitation. **The Lancet**, *377*(9778), 1693–1702. [[https://doi.org/10.1016/S0140-6736\(11\)60325-5](https://doi.org/10.1016/S0140-6736(11)60325-5)]([https://www.google.com/search?q=https://doi.org/10.1016/S0140-6736\(11\)60325-5](https://www.google.com/search?q=https://doi.org/10.1016/S0140-6736(11)60325-5))
 - 24) Langhorne, P., Coupar, F., & Pollock, A. (2009). Motor recovery after stroke: A systematic review. **The Lancet Neurology**, *8*(8), 741–754.
 - 25) Liepert, J., Bauder, H., Wolfgang, H. R., Miltner, W. H., Taub, E., & Weiller, C. (2000). Treatment-induced cortical reorganization after stroke in humans. **Stroke**, *31*(6), 1210–1216.
 - 26) Liepert, J., Uhde, I., Gräf, S., Leidner, O., & Weiller, C. (2001). Motor cortex plasticity during forced-use therapy in stroke patients: A preliminary study. **Journal of Neurology**, *248*(4), 315–321.
 - 27) Lyle, R. C. (1981). A performance test for assessment of upper limb function in physical rehabilitation treatment and research. **International Journal of Rehabilitation Research**, *4*(4), 483–492.
 - 28) Mark, V. W., Taub, E., & Morris, D. M. (2006). Neuroplasticity and constraint-induced movement therapy. **Eura Medicophys**, *42*(3), 269–280.
 - 29) Miltner, W. H., Bauder, H., Sommer, M., Dettmers, C., & Taub, E. (1999). Effects of constraint-induced movement therapy on chronic stroke patients: A replication. **Stroke**, *30*(3), 586–592.
 - 30) Morris, D. M., Taub, E., & Mark, V. W. (2006). Constraint-induced movement therapy: Characterizing the intervention protocol. **NeuroRehabilitation**, *21*(3), 167–185.
 - 31) Moxon, K. A., Oliviero, A., Aguilar, J., & Foffani, G. (2014). Cortical reorganization after spinal cord injury and stroke: Mechanisms and clinical implications. **Frontiers in Integrative Neuroscience**, *8*, Article 88.
 - 32) Nudo, R. J. (2006). Mechanisms for recovery of motor function following cortical damage. **Current Opinion in Neurobiology**, *16*(6), 638–644.
 - 33) Nudo, R. J., Wise, B. M., SiFuentes, F., & Milliken, G. W. (1996). Neural substrates for the effects of functional training on motor recovery after ischemic infarct. **Science**, *272*(5269), 1791–1794.
 - 34) Page, S. J., Sisto, S., Levine, P., & Johnston, M. V. (2002). Modified constraint-induced therapy: A randomized feasible and efficacy study. **Journal of Rehabilitation Research and Development**, *39*(5), 583–590.
 - 35) Page, S. J., Sisto, S. A., Levine, P., Johnston, M. V., & Hughes, M. (2001). Modified constraint-induced therapy: A randomized controlled trial. **Archives of Physical Medicine and Rehabilitation**, *82*(10), 1333–1339.

- 36) Page, S. J., Levine, P., Leonard, A., Szaflarski, J. P., & Kissela, B. M. (2008). Modified constraint-induced therapy in chronic stroke: Results of a single-blinded randomized controlled trial. **Physical Therapy**, *88*(3), 333–340.
- 37) Page, S. J., Levine, P., & Leonard, A. C. (2007). Modified constraint-induced therapy in acute stroke: A randomized controlled trial. **Neurorehabilitation and Neural Repair**, *21*(1), 50–58.
- 38) Pulse, A., & Jones, L. (2017). Dopaminergic pathways and reward-driven motor learning in neurorehabilitation. **Neuroscience & Biobehavioral Reviews**, *79*, 89–98.
- 39) Rotem-Kohavi, N., Oberlander, T. F., & Virji-Babul, N. (2014). BDNF expression and activity-dependent plastic changes in motor recovery: A systematic review. **Neurorehabilitation and Neural Repair**, *28*(8), 789–801.
- 40) Rothenberg, J., & Schmidt, R. (2019). Executive control and frontoparietal networks in post-stroke rehabilitation. **Cognitive, Affective, & Behavioral Neuroscience**, *19*(2), 201–215.
- 41) Rotter, J. B. (1966). Generalized expectancies for internal versus external control of reinforcement. **Psychological Monographs: General and Applied**, *80*(1), 1–28.
- 42) Smania, N., Aglioti, S. M., Cosentino, A., Camin, M., Gandolfi, M., Tinazzi, M., & Fiaschi, A. (2007). Application of constraint-induced movement therapy in patients with chronic stroke: A randomized controlled trial. **Journal of Rehabilitation Medicine**, *39*(8), 642–647.
- 43) Sterr, A., Elbert, T., Berthold, I., Kölbels, S., Rockstroh, B., & Taub, E. (2002). Longer term training in constraint-induced movement therapy in chronic hemiparesis. **Stroke**, *33*(3), 762–767.
- 44) Taub, E. (1980). Somatosensory deafferentation research with monkeys: Implications for rehabilitation medicine. In L. P. Ince (Ed.), **Behavioral psychology in rehabilitation medicine: Clinical applications** (pp. 371–401). Williams & Wilkins.
- 45) Taub, E., Crago, J. E., Burgio, L. D., Groomes, T. E., Cook, E. W., DeLuca, S. C., & Miller, N. E. (1994). An operant approach to rehabilitation medicine: Overcoming learned nonuse by shaping. **Journal of the Experimental Analysis of Behavior**, *61*(2), 281–293. <https://doi.org/10.1901/jeab.1994.61-281>
- 46) Taub, E., Miller, N. E., Novack, T. A., Cook, E. W., Fleming, W. C., Nepomuceno, C. S., Connell, J. S., & Crago, J. E. (1993). Technique to improve chronic motor deficit after stroke. **Archives of Physical Medicine and Rehabilitation**, *74*(4), 347–354.
- 47) Taub, E., Uswatte, G., & Pidikiti, R. (1999). Constraint-Induced Movement Therapy: A new family of techniques with broad application to physical rehabilitation—a clinical review. **Journal of Rehabilitation Research and Development**, *36*(3), 237–251.
- 48) Taub, E., Uswatte, G., Mark, V. W., & Morris, D. M. (2006). The transfer package: A essential component of Constraint-Induced Movement Therapy to improve daily arm use. **Physical Therapy**, *86*(11), 1548–1558.
- 49) van der Lee, J. H., Wagenaar, R. C., Lankhorst, G. J., Vogelaar, T. W., Devillé, W. L., & Bouter, L. M. (1999). Forced use of the upper extremity in chronic stroke patients: Results from a single-blind randomized clinical trial. **Stroke**, *30*(11), 2369–2375.
- 50) Veerbeek, J. M., van Wegen, E., van Peppen, R., van der Wees, P. J., Hendriks, E., Rietberg, M., & Kwakkel, G. (2014). What is the evidence for physical therapy poststroke? A systematic review and meta-analysis. **PLOS ONE**, *9*(2), e87987.
- 51) World Health Organization. (2001). **International classification of functioning, disability and health: ICF**. World Health Organization.
- 52) Wolf, S. L., Winstein, C. J., Miller, J. P., Taub, E., Uswatte, G., Morris, D., Giuliani, C., Light, K. E., & Nichols-Larsen, D. (2006). Effect of constraint-induced movement therapy on upper extremity function 3 to 9 months after stroke: The EXCITE randomized clinical trial. **JAMA**, *296*(17), 2095–2104. [<https://doi.org/10.1001/jama.296.17.2095>](<https://www.google.com/search?q=https://doi.org/10.1001/jama.296.17.2095>)
- 53) Wolf, S. L., Thompson, P. A., Winstein, C. J., Miller, J. P., Blanton, S. R., Croarkin, E., & EXCITE Investigators. (2010). The EXCITE stroke trial: Comparing early and delayed constraint-induced movement therapy. **Neurorehabilitation and Neural Repair**, *24*(3), 223–234.

- 54) Woytowicz, E. J., Rietschel, J. C., Goodman, R. N., Conroy, S. S., Sorkin, J. D., Whittall, J., & McCombe Waller, S. (2016). Determining levels of upper extremity impairment in stroke. *Journal of Neurologic Physical Therapy*, *40*(3), 184–192.
- 55) Zheng, Q. X., Ge, L., & Wang, C. (2019). The efficacy of modified constraint-induced movement therapy for upper limb function in stroke patients: A meta-analysis of randomized controlled trials. *Frontiers in Neurology*, *10*, Article 1029.
[<https://doi.org/10.3389/fneur.2019.01029>](<https://www.google.com/search?q=https://doi.org/10.3389/fneur.2019.01029>)