

Mitigating The Quick-Fix Fallacy and Kinesiophobia Following Orthobiologic Interventions: A Deterministic Health Informatics Architecture For Longitudinal Sports Rehabilitation

Pavan Manikanta Raghava Kasturi¹, Vithya Ganesan²

^{1,2} Department of Computer Science and Engineering,
Koneru Lakshmaiah Education Foundation, Vijayawada, Andhra Pradesh, India.
¹ kasturikpmr@gmail.com, ² vithyamtech@gmail.com
Corresponding Author: 1 Pavan Manikanta Raghava Kasturi
Email: kasturikpmr@gmail.com

ABSTRACT

Background: Regenerative orthobiologics, primarily Platelet-Rich Plasma (PRP) and Hyaluronic Acid (HA), induce rapid biochemical analgesia. In sports rehabilitation, this early pain reduction frequently triggers a "quick-fix fallacy," causing athletes to initiate premature loading before structural collagen maturation. Concurrently, kinesiophobia impedes necessary mechanotransduction during episodic follow-up care.

Objective: To design, implement, and computationally benchmark a deterministic assistive health informatics architecture that continuously bridges the outpatient monitoring gap, preventing premature tissue overload and mitigating kinesiophobia.

Methods: A three-tiered clinical decision support framework was constructed. Tier-1 captures asynchronous Ecological Momentary Assessments (EMA) measuring Visual Analog Scale (VAS) pain, Rating of Perceived Exertion (RPE), and Tampa Scale for Kinesiophobia (TSK-11) on a mobile edge node with hash-indexed storage ($O(1)$ access). Tier-2 enforces HL7 FHIR Release 4 JSON serialisation over TLS 1.3 and AES-256-GCM encryption. Tier-3 executes a deterministic $O(1)$ boolean rule engine evaluated against clinical healing envelopes under Human-in-the-Loop (HITL) physician governance.

Results: In empirical verification benchmarks on 10,000 synthetic patient trajectories, threshold evaluation latency averaged 0.0006 ± 0.0002 ms ($0.59 \mu\text{s}$). Rapid analgesia ($\Delta\text{VAS} \leq -50\%$ within 14 days) combined with excessive loading ($\text{RPE} > \text{RPE}_{\text{max_safe}}$) triggered an OVERLOAD_WARNING. Low pain ($\text{VAS} \leq 3$) with rising fear-avoidance ($\Delta\text{TSK-11} > +10\%$ after Day 21) reliably triggered a PHOBIC_ARREST triage alert.

Conclusion: This deterministic architecture eliminates outpatient monitoring blind spots, providing clinicians with real-time, explainable telemetry to protect collagen remodelling and resolve kinesiophobia.

Keywords: Platelet-Rich Plasma, Hyaluronic Acid, Sports Medicine, Orthopaedic Rehabilitation, Kinesiophobia, Clinical Decision Support Systems.

1. INTRODUCTION

Sports traumatology and orthopaedics have shifted from palliative symptom management toward regenerative biological repair through intra-articular and intra-tendinous orthobiologics, primarily Platelet-Rich Plasma (PRP) and Hyaluronic Acid (HA) [1]. While these biological agents demonstrate clinical efficacy in resolving focal chondral lesions, midsubstance ligamentous sprains, and chronic tendinopathies, real-world therapeutic outcomes exhibit substantial heterogeneity [1]. Clinical failure frequently stems from a systemic post-procedure infrastructural limitation: the inability to continuously monitor and modulate the athlete's high-frequency psychobiological recovery trajectory during outpatient rehabilitation.

The conventional outpatient delivery model relies on episodic, retrospective clinical consultations conducted at arbitrary 3- to 6-week intervals. This structure creates an unmonitored "temporal gap" between hospital visits, blinding orthopaedic surgeons and sports physiotherapists to the patient's daily mechanical loading events, fluctuating nociceptive baselines, and evolving psychological states. Within this unmonitored interval, two diametrically opposed clinical failure modes emerge: biomechanical tissue overload (the "quick-fix fallacy") and neuro-cognitive immobilisation (kinesiophobia).

1.1. Biochemical Analgesia versus Structural Maturation

The quick-fix fallacy originates from the temporal divergence between molecular biochemical anti-inflammation and mechanical extracellular matrix (ECM) remodelling. Autologous PRP preparations deliver concentrated platelet α -granules containing bioactive cytokines and growth factors, including Transforming Growth Factor-Beta 1 (TGF- β 1), Vascular Endothelial Growth Factor (VEGF), Platelet-Derived Growth Factor (PDGF), and basic Fibroblast Growth Factor (bFGF) [1]. In parallel, high-molecular-weight HA provides immediate viscosupplementation while binding cluster determinant 44 (CD44) and the receptor for hyaluronan-mediated motility (RHAMM) on synovial fibroblasts and articular chondrocytes [2]. This cascade downregulates pro-

inflammatory cytokines, specifically Interleukin-1 β (IL-1 β) and Tumour Necrosis Factor- α (TNF- α), suppressing intra-articular nociceptive signalling and delivering substantial analgesia within 7 to 14 days post-injection [2]. However, extracellular matrix remodelling, specifically the biological transition from disorganised, hypercellular Type III collagen to high-tensile, linearly aligned Type I collagen bundles mediated by lysyl oxidase cross-linking, requires between 12 and 24 weeks of protected, progressive mechanotransduction [3],[4]. Interpreting early clinical analgesia as complete tissue restitution, athletes prematurely resume unauthorised high-velocity, ballistic, or plyometric loading. This excessive stress exceeds the ultimate tensile yield strength of nascent collagen fibrils, causing micro-structural disruption, biological graft attenuation, or complete tendinopathic recurrence [3]. These failure dynamics are particularly prevalent in patellar tendinopathy (jumper's knee), midportion Achilles tendinopathy, lateral epicondylalgia, and focal femoral condyle chondral lesions.

1.2. Neurophysiological Fear-Avoidance and Kinesiophobia

At the opposite end of the spectrum, functional rehabilitation is frequently compromised by maladaptive neuro-cognitive fear-avoidance behaviours, as conceptualised in the Fear-Avoidance Model [5]. Following acute structural trauma, nociceptive signalling through primary A- δ and unmyelinated C fibres triggers neuroplastic adaptations within the somatosensory cortex, anterior cingulate cortex, and amygdala [5]. In athletes with heightened anxiety, the perceived vulnerability of the repaired structure outlasts the initial nociceptive stimulus. This psychological state manifests as kinesiophobia, an excessive, debilitating fear of movement and reinjury [6]. Empirical meta-analyses indicate that nearly 45% of athletes fail to return to their pre-injury competitive baseline despite fully intact, biologically healed tissue architectures [7]. Neurophysiologically, kinesiophobia disrupts standard motor unit recruitment patterns, inducing abnormal muscle co-contraction, compensatory biomechanical loading, and arthrogenic muscle inhibition [6].

Because connective tissue remodelling depends fundamentally on mechanotherapy, wherein controlled strain stimulates cellular integrins to initiate intracellular signalling pathways that upregulate collagen transcription [4], fear-induced immobilisation deprives the healing matrix of the mechanical cues essential for alignment. This results in disuse atrophy, capsular contracture, and joint stiffness. While psychometric questionnaires such as the 11-item Tampa Scale for Kinesiophobia (TSK-11) can quantify fear-avoidance [8], standard episodic administration misses these acute behavioural inflections.

1.3. Translational Clinical Relevance

- **What is already known:** Intra-articular and intra-tendinous orthobiologics induce rapid localised anti-inflammatory analgesia. However, standard post-injection outpatient care remains episodic (intermittent 3- to 6-week reviews), relying on retrospective patient recall that is vulnerable to cognitive bias.
- **What this study adds:** This work establishes a deterministic, real-time health informatics architecture that continuously tracks daily psychobiological trajectories, actively preventing premature tendon/graft overload while simultaneously identifying kinesiophobia-induced functional stagnation without relying on opaque, black-box artificial intelligence models.

2. METHODS

2.1. Study Setting and Design

This study establishes a system architecture and computational verification framework for outpatient sports medicine clinics, academic orthopaedic departments, and athletic training centres. To prevent diagnostic liability, the framework formalises deterministic data pipelines, cryptographic security, and rule-based state logic prior to longitudinal clinical trials. Table 1 contrasts this architecture with conventional care.

Table 1: Episodic vs. Deterministic Continuous Monitoring Paradigms

System Parameter	Traditional Outpatient Rehabilitation	Proposed Deterministic EMA Architecture
Data Acquisition Frequency	Episodic (Every 3 to 6 weeks)	Continuous (Asynchronous daily polling)
Data Integrity	Vulnerable to recall bias and peak-end rule	High fidelity (Ecological momentary capture)
Interoperability Standards	Proprietary databases or unindexed PDFs	Standardised HL7 FHIR Release 4 serialisation

Clinical Intervention Point	Reactive (Damage established before visit)	Proactive (Real-time anomaly triage notifications)
Computational Governance	Unstructured manual chart review	Deterministic O(1) HITL rule engine (No black-box AI)

Table Note: Abbreviations: EMA: Ecological Momentary Assessment, FHIR: Fast Healthcare Interoperability Resources, HITL: Human-in-the-Loop, HL7: Health Level Seven.

2.2. Multi-Tiered System Architecture

The framework is structured into three operational layers, summarised in Table 2.

Table 2: System Architecture and Data Processing Flow

Architectural Tier	Sub-Component	Technical Mechanism / Data Flow
Tier-1: Patient Edge Node	Asynchronous EMA Polling	Capture of VAS (0 to 10), Borg CR10 RPE (0 to 10), and TSK-11 (11 to 44)
	Local Storage Engine	SQLite transactional cache utilising hash indexing (O(1) access)
Tier-2: Secure Transmission	Data Normalisation	Structural serialisation into HL7 FHIR Release 4 JSON resources
	Transport and Storage Security	End-to-end TLS 1.3 transport to AES-256-GCM rest encryption
Tier-3: Clinical Decision Support	Deterministic State Engine	Evaluates continuous telemetry against therapeutic envelopes
	Silent Execution Path	Automated time-series logging into Electronic Medical Record (EMR)
	Alert Trigger Path	Generates high-priority OVERLOAD_WARNING or PHOBIC_ARREST flags
	Human-in-the-Loop (HITL)	Clinical review, diagnostic validation, and intervention by physician

Table Note: Abbreviations: AES: Advanced Encryption Standard, EMA: Ecological Momentary Assessment, EMR: Electronic Medical Record, GCM: Galois/Counter Mode, HITL: Human-in-the-Loop, JSON: JavaScript Object Notation, RPE: Rating of Perceived Exertion, TLS: Transport Layer Security, TSK-11: Tampa Scale for Kinesiophobia-11, VAS: Visual Analog Scale.

2.3. Tier-1: Patient Edge Node and Asynchronous EMA

Tier-1 captures real-time recovery data directly within the patient's daily natural environment via a mobile application (Edge Node) [9]. An asynchronous, pseudo-randomised notification schedule within configured waking hours prevents anticipatory behavioural skewing. Smartphone-based EMA methodologies have demonstrated high compliance and construct validity for longitudinal musculoskeletal monitoring [10],[11].

The application records:

1. **Nociceptive Index:** Visual Analog Scale (VAS, 0 to 10) measuring localised anatomical pain.
2. **Biomechanical Exertion Index:** Modified Borg CR10 Rating of Perceived Exertion (RPE, 0 to 10) capturing physical session intensity and unintended mechanical overload.
3. **Psychometric Fear-Avoidance Index:** Digitised TSK-11 (score range 11 to 44) evaluating movement-related fear and somatic focus [8].

The data schema also accommodates disease-specific functional tools, such as VISA-P, VISA-A, and IKDC forms. An edge-side SQLite database structures records as key-value pairs indexed by UTC timestamps. This configuration achieves O(1) constant time complexity for data insertion and retrieval, preserving battery life and securing offline data during network interruptions.

2.4. Tier-2: HL7 FHIR Serialisation and Cryptographic Security

Tier-2 standardises data using Health Level Seven Fast Healthcare Interoperability Resources (HL7 FHIR) Release 4 JSON objects [12]. Metrics are mapped to standardised vocabularies:

- VAS pain scores: LOINC code 72514-3 (Pain severity - 0-10 verbal numeric rating).
- RPE loading values: LOINC code 72216-5 (Rating of perceived exertion).
- TSK-11 scores: LOINC code 90779-0 (Tampa Scale for Kinesiophobia-11).

Payloads include ISO 8601 UTC timestamps, patient UUIDs, and calibrated device metadata. Network transmission mandates TLS 1.3 with Ephemeral Elliptic Curve Diffie-Hellman (ECDHE) key exchange for Perfect

Forward Secrecy. Data at rest is encrypted with symmetric AES-256-GCM (Authenticated Encryption with Associated Data), ensuring confidentiality and immediate rejection of altered packets.

3. RESULTS

3.1. Tier-3: Deterministic Rule Engine

Tier-3 serves as the clinical decision support core. While deep neural networks offer predictive power, their epistemic opacity ("black-box" nature) creates regulatory and clinical liabilities in acute care [13].

Tier-3 implements a rule-based deterministic finite state machine (FSM). Hard-coded boolean logic gates evaluate telemetry against physician-configured healing envelopes. Evaluating scalar metrics over bounded ranges yields $O(1)$ constant time complexity, maintaining instantaneous execution across large EMR repositories.

3.2. Algorithmic State Evaluation and Calibration

The engine evaluates longitudinal telemetry across time (t), pain (Δ VAS), strain (RPE), and fear-avoidance (Δ TSK-11). Parameters are detailed in Table 3.

Table 3: Tier-3 Algorithmic State Evaluation Parameters

Clinical State	Temporal Criteria (t)	Nociceptive Parameter (VAS)	Biomechanical Loading (RPE)	Psychometric State (TSK-11)	System Action / Triage Flag
Biomechanical Overload (Quick-Fix Fallacy)	$t < 14$ days	Δ VAS $\leq -50\%$ (Rapid drop)	$RPE > RPE_{\max_safe}$	Non-contributory	OVERLOAD_WARNING
Phobic Arrest (Kinesiophobia)	$t > 21$ days	VAS ≤ 3 (Controlled pain)	$RPE < RPE_{\min_required}$	Δ TSK-11 $> +10\%$ (Rising fear)	PHOBIC_ARREST
Normal Physiological Remodelling	$14 \leq t \leq 180$ days	Stable or gradual decline	$RPE_{\min} \leq RPE \leq RPE_{\max}$	Stable or progressive decline	LOG_DATA_ONLY

Table Note: Abbreviations: Δ TSK-11: Percentage change in Tampa Scale for Kinesiophobia, Δ VAS: Percentage change in Visual Analog Scale, $RPE_{\max_safe}$: Clinician-defined maximum safe Rating of Perceived Exertion, $RPE_{\min_required}$: Minimum exertion required for mechanotransductive remodelling.

The mathematical evaluation pathways operate as follows:

$$\Delta$$
VAS = $\left(\frac{VAS_{\text{current}} - VAS_{\text{baseline}}}{VAS_{\text{baseline}}} \right) \times 100$

$$\Delta$$
TSK-11 = $\left(\frac{TSK-11_{\text{current}} - TSK-11_{\text{baseline}}}{TSK-11_{\text{baseline}}} \right) \times 100$

The deterministic engine supports customised threshold matrices tailored to specific anatomical healing kinetics:

- **Patellar and Achilles Tendinopathies:** Strict initial limits ($RPE_{\max_safe} \leq 3$) during the 14-day inflammatory phase to prevent tensile yield failure, transitioning to mandatory eccentric mechanotransduction ($RPE_{\min_required} \geq 4$) by Day 28.
- **Focal Chondral Lesions:** Extended protection ($t < 21$ days, $RPE_{\max_safe} \leq 2$) to safeguard early chondrocyte adherence and viscosupplementation integration before permitting closed-kinetic-chain axial loading.
- **Lateral Epicondylalgia:** Calibrated against repetitive wrist extensor and gripping strain spikes.

Three operational states execute deterministically:

- **State 1: Biomechanical Overload (OVERLOAD_WARNING):** In acute recovery ($t < 14$ days), if rapid analgesia (Δ VAS $\leq -50\%$) coincides with high loading ($RPE > RPE_{\max_safe}$), the engine flags early Type III collagen disruption and alerts the clinician.

- **State 2: Phobic Arrest (PHOBIC_ARREST):** In subacute remodelling ($t > 21$ days), if pain is low ($VAS \leq 3$) but fear rises ($\Delta TSK-11 > +10\%$) alongside deficient loading ($RPE < RPE_{min_required}$), the system alerts the team to initiate cognitive reassurance and progressive loading.
- **State 3: Physiological Remodelling (LOG_DATA_ONLY):** If values remain within bounds, telemetry is indexed silently into the EMR [14].

3.3. Human-in-the-Loop Clinical Governance

The architecture enforces strict Human-in-the-Loop (HITL) governance [15]. The engine possesses zero autonomous prescription authority. When state boundaries are breached, it generates an encrypted HL7 FHIR Flag resource sent to the orthopaedic surgeon's dashboard. The clinician evaluates the telemetry and determines appropriate clinical action [15].

3.4. Simulated Clinical Scenarios

The deterministic engine was verified using two representative recovery trajectories:

- **Scenario A (Overload Divergence):** An athlete receiving PRP for patellar tendinopathy logs baseline $VAS = 7$ and $TSK-11 = 28$. On Day 8, pain drops to $VAS = 2$ ($\Delta VAS = -71.4\%$). The patient performs unprescribed plyometric drills, logging $RPE = 8$ (configured $RPE_{max_safe} = 4$). In constant time ($O(1)$), the engine identifies the threshold breach, triggers an OVERLOAD_WARNING, and alerts the sports medicine team.
- **Scenario B (Phobic Arrest Divergence):** An athlete receiving HA for a Grade II chondral defect logs baseline $VAS = 6$ and $TSK-11 = 24$. By Day 28, pain drops to $VAS = 2$. Due to reinjury anxiety, the patient avoids loading, recording $RPE = 1$ (configured $RPE_{min_required} = 4$) while fear scores rise to $TSK-11 = 34$ ($\Delta TSK-11 = +41.7\%$). The engine triggers a PHOBIC_ARREST alert, prompting the team to restore progressive mechanotransduction.

3.5. Computational Performance Benchmarks

Operational efficiency was empirically benchmarked across 10,000 synthetic patient trajectories (representing 180 daily telemetry entries per patient, totalling 1.8 million evaluations):

- **Evaluation Latency:** Mean execution time per payload was 0.0006 ± 0.0002 ms (sub-microsecond scale), validating strictly bounded $O(1)$ constant time complexity suitable for high-throughput hospital EMR integration without processing bottlenecks.
- **Edge Storage Footprint:** SQLite transactional cache consumption on the mobile client was measured at 16.00 KB for a complete 180-day trajectory (≈ 89 bytes per daily entry including table indexing and database page allocation overhead), imposing negligible memory overhead on edge hardware.
- **Network Overhead:** Standardised HL7 FHIR Release 4 JSON payload size was measured at 0.50 KB (512 bytes) per multi-variable observation, supporting instantaneous asynchronous transmission across low-bandwidth or intermittent mobile network connections.

3.6. Kinematic Sensor Integration and Translational Roadmap

While client-side mobile EMA provides high-fidelity subjective metrics, future translational iterations can incorporate passive kinematic biosensing arrays to complement subjective RPE inputs.

Integrating wearable Inertial Measurement Units (IMUs) equipped with tri-axial accelerometers and gyroscopes secured adjacent to the treated anatomical site enables continuous, objective quantification of mechanical load. Kinematic parameters such as peak ground reaction forces, joint range-of-motion angular velocity, and limb-loading asymmetry indices can be sampled continuously.

To prevent data transmission bottlenecks, edge-computing algorithms on the mobile device can execute real-time data decimation and statistical feature extraction, translating continuous high-frequency kinematic streams into discrete daily mechanical work indices. As illustrated in the end-to-end data pipeline (Figure 1) and the underlying state logic (Figure 2), these indices feed directly into the Tier-3 engine's threshold comparisons, corroborating self-reported RPE exertion against physical limb mechanics.




Figure 1. Three-tiered health informatics architecture for continuous post-orthobiologic monitoring. Schematic illustrating Tier-1 edge data capture (VAS, RPE, TSK-11), Tier-2 HL7 FHIR Release 4 JSON serialisation and TLS 1.3/AES-256-GCM encryption, and Tier-3 deterministic O(1) clinical decision support interfacing directly with hospital EMR systems under Human-in-the-Loop governance.




Figure 2. Deterministic finite state machine logic and alert branching.

Flowchart of the O(1) scalar threshold evaluation algorithm illustrating the temporal, nociceptive, mechanical, and psychometric decision gates that separate physiological collagen remodelling from acute overload and phobic arrest states.

4. DISCUSSION

This three-tiered health informatics architecture addresses the post-orthobiologic monitoring gap by translating continuous psychobiological telemetry into actionable, explainable clinical alerts.

4.1. Divergence from Conventional Patient Portals

Standard commercial patient portals operate as passive survey repositories. They capture data at fixed intervals without evaluating cross-variable relationships. If an athlete experiences pain relief and subsequently overloads healing tissue, standard portals log the low pain score as an uncontextualized improvement. In contrast, this closed-loop triage engine cross-references nociception drops against strain spikes, identifying hidden risks before structural failure occurs.

4.2. Implementation and Interoperability

While Tier-2 uses HL7 FHIR Release 4 JSON serialisation [12], many clinical systems rely on legacy HL7 v2 messaging or siloed relational databases. In such environments, lightweight API translation gateways or FHIR-facade adapters parse incoming payloads into standard EHR views, enabling cross-institutional integration.

4.3. Patient Compliance

Long-term adherence is critical in mobile health deployments [9],[11]. To reduce prompt fatigue, the mobile interface uses micro-surveys taking under 30 seconds to complete, combined with adaptive sampling that reduces query frequency once psychological stability is established.

4.4. Mitigating Alert Fatigue and Clinical Liability

Alert fatigue is a common cause of decision-support failure [14]. The architecture minimises false positives by avoiding uncalibrated thresholds. Surgeons customise threshold envelopes based on individual tissue biology, distinguishing elite athletes from recreational patients. Enforcing strict boolean logic gates (Time $\cap$ Pain $\cap$ Load $\cap$ Psychology) suppresses baseline data and elevates only actionable anomalies [14],[15], reducing chart-review burden while documenting objective compliance history in the EMR.

5. CONCLUSION

Regenerative orthobiologics produce rapid biochemical analgesia that requires modernised post-injection monitoring. Conventional episodic follow-up leaves athletes vulnerable to premature mechanical failure or kinesiophobia-induced recovery arrest.

The deterministic health informatics architecture presented here resolves this longitudinal monitoring deficit. Combining edge-based asynchronous EMA data collection, HL7 FHIR Release 4 interoperability, and

cryptographic security with an $O(1)$ boolean rule engine under strict Human-in-the-Loop governance, the platform delivers transparent clinical triage. This framework protects collagen remodelling, supports objective mechanobiology, and optimises functional sports rehabilitation.

REFERENCES

1. Filardo G, Di Matteo B, Kon E, Merli G, Marcacci M. Platelet-rich plasma in tendon-related disorders: results and indications. *Knee Surg Sports Traumatol Arthrosc.* 2018 Jul;26(7):1984–1999. doi: 10.1007/s00167-016-4261-4.
2. Altman RD, Manjoo A, Fierlinger A, Niazi F, Nicholls M. The mechanism of action for hyaluronic acid treatment in the osteoarthritic knee: a systematic review. *BMC Musculoskelet Disord.* 2015 Oct 26;16:321. doi: 10.1186/s12891-015-0775-z.
3. Sharma P, Maffulli N. Biology of tendon injury: healing, modeling and remodeling. *J Musculoskelet Neuronal Interact.* 2006 Apr-Jun;6(2):181-190. PMID: 16849830.
4. Khan KM, Scott A. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *British Journal of Sports Med.* 2009 Apr;43(4):247-252. doi: 10.1136/bjsm.2008.054239.
5. Vlaeyen JW, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain.* 2000 Apr;85(3):317-332. doi: 10.1016/S0304-3959(99)00242-0.
6. Kvist J, Ek A, Sporrstedt K, Good L. Fear of re-injury: a hindrance for returning to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2005 Jul;13(5):393–397. doi: 10.1007/s00167-004-0591-8.
7. Ardern CL, Taylor NF, Feller JA, Webster KE. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *British Journal of Sports Med.* 2014 Nov;48(21):1543-1552. doi: 10.1136/bjsports-2013-093398.
8. Woby SR, Roach NK, Urmston M, Watson PJ. Psychometric properties of the TSK-11: a shortened version of the Tampa Scale for Kinesiophobia. *Pain.* 2005 Sep;117(1-2):137-144. doi: 10.1016/j.pain.2005.05.029.
9. Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annu Rev Clin Psychol.* 2008 Apr;4:1-32. doi: 10.1146/annurev.clinpsy.3.022806.091415.
10. Kaur E, Haghighi PD, Cicuttini FM, Urquhart DM. Smartphone-Based Ecological Momentary Assessment for Collecting Pain and Function Data for Those with Low Back Pain. *Sensors (Basel)* 2022 Sep;22(18):7095. doi:10.3390/s22187095.
11. Moore RC, Swendsen J, Depp CA. Applications for self-administered mobile cognitive assessments in clinical research: A systematic review. *Int J Methods Psychiatr Res.* 2017 Dec;26(4):e1562. doi: 10.1002/mpr.1562.
12. Bender D, Sartipi K. HL7 FHIR: An Agile and RESTful approach to healthcare information exchange. In: *Proceedings of the 26th IEEE International Symposium on Computer-Based Medical Systems.* York (PA): IEEE; 2013. 326-331. doi: 10.1109/CBMS.2013.6627810.
13. Holzinger A, Langs G, Denk H, Zatloukal K, Müller H. Causability and explainability of artificial intelligence in medicine. *WIREs Data Mining Knowl Discov.* 2019 Jul-Aug;9(4):e1312. doi: 10.1002/widm.1312.
14. Sutton RT, Pincock D, Baumgart DC, Sadowski DC, Fedorak RN, Kroeker KI. An overview of clinical decision support systems: benefits, risks, and strategies for success. *NPJ Digit Med.* 2020 Feb 6;3:17. doi: 10.1038/s41746-020-0221-y.
15. Holzinger A. Interactive machine learning for health informatics: when do we need the human-in-the-loop? *Brain Inform.* 2016 Mar;3:119-131. doi: 10.1007/s40708-016-0042-6.